Reference number(s)
4201-A

Standard Medicare Part B Management Kanuma

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Kanuma	sebelipase alfa

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications[1]

Kanuma is indicated for the treatment of patients with a diagnosis of Lysosomal Acid Lipase (LAL) deficiency.

All other indications will be assessed on an individual basis. Submissions for indications other than those listed in this criteria should be accompanied by supporting evidence from Medicare approved compendia.

Documentation

The following documentation must be available, upon request, for all submissions:

- For initial requests: lysosomal acid lipase enzyme assay or genetic testing results supporting diagnosis.
- Continuation requests: lab values or chart notes documenting benefit from therapy.

© 2025 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Reference number(s)
4201-A

Coverage Criteria

Lysosomal Acid Lipase (LAL) Deficiency[1]

Authorization of 12 months may be granted for treatment of LAL deficiency when both of the following criteria are met:

- Diagnosis of LAL deficiency was confirmed by enzyme assay demonstrating a deficiency of lysosomal acid lipase enzyme activity or by genetic testing; AND
- Member has alanine aminotransferase level (ALT) greater than or equal to 1.5 times the upper limit of normal (based on the age and gender-specific normal ranges) on two consecutive ALT measurements obtained at least one week apart.

Continuation of Therapy

All members (including new members) requesting authorization for continuation of therapy must be currently receiving therapy with the requested agent.

Authorization of 12 months may be granted when all of the following criteria are met:

- The member is currently receiving therapy with the requested medication.
- The requested medication is being used to treat an indication in the coverage criteria.
- The member is receiving benefit from therapy (e.g., improvement, stabilization, or slowing of disease progression for weight-for-age z-score if exhibiting growth failure, low-density lipoprotein [LDL], high-density lipoprotein [HDL], triglycerides, or alanine aminotransferase [ALT]).

Summary of Evidence

The contents of this policy were created after examining the following resources:

- The prescribing information for Kanuma.
- The available compendium
    - National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium
    - Micromedex DrugDex
    - American Hospital Formulary Service- Drug Information (AHFS-DI)
    - Lexi-Drugs
    - Clinical Pharmacology
- Lysosomal Acid Lipase Deficiency (Endotext).

After reviewing the information in the above resources, the FDA-approved indications listed in the prescribing information for Kanuma are covered.

© 2025 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Reference number(s)
4201-A

Explanation of Rationale

Support for FDA-approved indications can be found in the manufacturer's prescribing information.

Support for confirming the diagnosis with enzyme assays or genetic testing prior to using Kanuma to treat LAL deficiency can be found in an article by Wilson and Patni. LAL deficiency can be diagnosed by demonstrating deficient LAL enzyme activity, as well as by genetic testing identifying pathogenic variants of the LIPA gene. Historically, enzyme activity was measured in cultured fibroblasts, peripheral leukocytes, or liver tissue. A newer method has been developed to determine LAL activity. This method measures LAL activity in dried blood spots (DBS), and uses Lalistat 2, a highly specific inhibitor of LAL. LAL activity is determined by comparing total lipase activity to lipase activity with Lalistat 2.

References

1. Kanuma [package insert]. Cheshire, CT: Alexion Pharmaceuticals Inc.; July 2024.
2. Wilson DP, Patni N. Lysosomal Acid Lipase Deficiency. [Updated 2023 Mar 15]. In: Feingold KR, Anawalt B, Blackman MR, et al., editors. Endotext [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK395569/

© 2025 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.